

TESTOPEL[®] 75mg
(testosterone pellets) 

TESTOSTERONE REPLACEMENT THERAPY

TURN IT



**One dose of TESTOPEL[®] can normalize
T-levels for 3-4 months.**

866.SLATE.50 • www.testopel.com

IMPORTANT SAFETY INFORMATION

TESTOPEL is indicated for replacement therapy in conditions associated with low or no endogenous testosterone.




TESTOPEL should not be used in men with breast cancer due to the potential for development of hypercalcemia or in men with prostate cancer due to increased risk for the development of prostatic carcinoma. Patients should not use TESTOPEL if they have had a previous adverse reaction to TESTOPEL. TESTOPEL is not approved for use in women and may cause fetal harm.

**See enclosed Important Safety Information and complete
Prescribing Information for TESTOPEL.**

TESTOPEL[®] (testosterone pellets)

TESTOPEL is the only FDA-approved implantable testosterone pellet. Once placed, **TESTOPEL** most commonly normalizes testosterone levels for 3-4 months. **TESTOPEL** placement is done in the doctor's office 2-4 times a year.

For busy men taking medication every day can be a real hassle. Skipping or missing doses may cause a significant drop in testosterone level, possibly resulting in the re-emergence of symptoms associated with low testosterone.

TREATMENT OPTIONS	 TESTOPEL[®]	 Topical Gels	 Injection
DOSING	Every 3-6 months	Daily	Every 2-4 weeks

With **TESTOPEL** you have the confidence of knowing your testosterone replacement medication is available to meet your body's demands for 3-4 months per dose. In fact, **TESTOPEL** is the only long-acting testosterone pellet approved for use by the FDA.

IMPORTANT SAFETY INFORMATION

Side effects reported with use of TESTOPEL include: excessive frequency and duration of penile erections, hirsutism, increased serum cholesterol, acne, acceleration of bone maturation without compensatory gain in linear growth in children, male pattern baldness, alterations in liver function tests, suppression of clotting factors, polycythemia, increased or decreased libido, headache, anxiety, depression, and generalized paresthesia. Some men may have breast development, breast discomfort, edema, or prostate enlargement accompanied by difficulty urinating.

Convenience.

Using **TESTOPEL** may help eliminate the following hassles:


- The need to take or apply a daily testosterone replacement medication.
- Concerns over accidental testosterone medication transfer to another person resulting from inadequate hand washing or contact with unwashed or unclothed areas of testosterone gel application.
- Monthly pharmacy visits and co-pays.
- The need to pack testosterone replacement medication when traveling.

TURN ON

Cost Savings.

- A national audit of over 35,000 insurance benefit confirmations reveals that 97% of **TESTOPEL** claims are reimbursed by insurance companies. For 50% of claims there is no patient out-of-pocket medication expense. The average patient out-of-pocket expense for the remaining claims is \$31.48.*

**Average Out-of-Pocket Expense
for Covered Claims**



\$31.48

\$0

**This is not a guarantee of insurance coverage or payment amount.*

Convenience.

Using **TESTOPEL** may help eliminate the following hassles:

- The need to take or apply a daily testosterone replacement medication.
- Concerns over accidental testosterone medication transfer to another person resulting from inadequate hand washing or contact with unwashed or unclothed areas of testosterone gel application.
- Monthly pharmacy visits and co-pays.
- The need to pack testosterone replacement medication when traveling.

TURN ON

Cost Savings.

- A national audit of over 35,000 insurance benefit confirmations reveals that 97% of **TESTOPEL** claims are reimbursed by insurance companies. For 50% of claims there is no patient out-of-pocket medication expense. The average patient out-of-pocket expense for the remaining claims is \$31.48.*

**Average Out-of-Pocket Expense
for Covered Claims**



**This is not a guarantee of insurance coverage or payment amount.*

TESTOPEL[®] (testosterone pellets)

TAKE THE LOW T QUIZ

- | | | | |
|------------------------------|-----------------------------|-----|---|
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | 1. | Do you have a decrease in libido (sex drive)? |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | 2. | Do you have a lack of energy? |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | 3. | Do you have a decrease in strength and/or endurance? |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | 4. | Have you lost height? |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | 5. | Have you noticed a decreased enjoyment in life? |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | 6. | Are you sad and/or grumpy? |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | 7. | Are your erections less strong? |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | 8. | Have you noted a recent deterioration in your ability to play sports? |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | 9. | Are you falling asleep after dinner? |
| <input type="checkbox"/> Yes | <input type="checkbox"/> No | 10. | Has there been a recent deterioration in your work performance? |

Yes No

Did you answer YES to either question 1 OR 7?

Yes No

Did you answer YES to at least three questions?

If you answered yes to questions 1 OR 7, or answered yes to any three questions, you may be suffering from Low T. Share your results with your doctor and ask about a simple blood test that can help better diagnose Low T and your treatment options.

Adapted from: Morley JE, Charlton E, Patrick P, et al. Validation of a screening questionnaire for androgen deficiency in aging males. *Metabolism*. 2000;49(9):1239-1242.

COMMONLY ASKED QUESTIONS ABOUT TESTOPEL®



What size is a TESTOPEL pellet?



Each **TESTOPEL** pellet is small.

The photo to the right shows the size of **TESTOPEL** relative to a dime.



Where is TESTOPEL placed?



TESTOPEL is placed just under the skin in the hip-area. Imagine the outer corner of the top of your pants back pocket.



How will my doctor know when to replace my TESTOPEL?



Your doctor may either schedule a follow-up visit or if you feel your symptoms returning you should contact your doctor so your testosterone levels can be measured.



What can I expect after my TESTOPEL placement?



You can return to work immediately. It is advisable not to swim, use a hot tub or lift heavy weights for a few days after **TESTOPEL** is placed, but you may otherwise resume normal activity. You may experience soreness and/or bruising at the site. Please follow your doctor's instructions.



Will my insurance pay for TESTOPEL?



The majority of insurance plans will cover **TESTOPEL**. It can be treated as either a medical or pharmacy benefit depending on your plan. In either case, you won't need to go to the pharmacy to get a **TESTOPEL** prescription filled. Your doctor's office can work with the **TESTOPEL** Reimbursement Program to determine your exact benefit and provide appropriate direction.

IMPORTANT SAFETY INFORMATION

TESTOPEL is indicated for replacement therapy in conditions associated with low or no endogenous testosterone.

TESTOPEL should not be used in men with breast cancer due to the potential for development of hypercalcemia or in men with prostate cancer due to increased risk for the development of prostatic carcinoma. Patients should not use TESTOPEL if they have had a previous adverse reaction to TESTOPEL. TESTOPEL is not approved for use in women and may cause fetal harm.

Side effects reported with use of TESTOPEL include: excessive frequency and duration of penile erections, hirsutism, increased serum cholesterol, acne, accelera-

tion of bone maturation without compensatory gain in linear growth in children, male pattern baldness, alterations in liver function tests, suppression of clotting factors, polycythemia, increased or decreased libido, headache, anxiety, depression, and generalized paresthesia. Some men may have breast development, breast discomfort, edema, or prostate enlargement accompanied by difficulty urinating.

TESTOPEL insertion may cause pain at the site of subcutaneous implantation of pellets and is rarely associated with anaphylactoid reactions. There is less flexibility for dosage adjustment compared to oral administration or intramuscular injection of oil solutions or aqueous suspensions. Surgical removal may be required if testosterone therapy is discontinued.